The information you received in the MPTC *Unload Your Backpack* training does not take the place of professional medical advice. These assessments, also available online, are for your informational and educational purposes. For questions or more information, consult with your primary care physician.

PCL-5

Instructions: Below is a list of problems that people sometimes have in response to a very stressful experience. Please read each problem carefully and then circle one of the numbers to the right to indicate how much you have been bothered by that problem in the past month.

In the past month, how much were you bothered by:	*Not at all*	*A little bit*	*Moderately*	*Quite a bit*	*Extremely*
1. Repeated, disturbing, and unwanted memories of the stressful experience?	0	1	2	3	4
2. Repeated, disturbing dreams of the stressful experience?	0	1	2	3	4
3. Suddenly feeling or acting as if the stressful experience were actually happening again *(as if you were actually back there reliving it)*?	0	1	2	3	4
4. Feeling very upset when something reminded you of the stressful experience?	0	1	2	3	4
5. Having strong physical reactions when something reminded you of the stressful experience *(for example, heart pounding, trouble breathing, sweating)*?	0	1	2	3	4
6. Avoiding memories, thoughts, or feelings related to the stressful experience?	0	1	2	3	4
7. Avoiding external reminders of the stressful experience *(for example, people, places, conversations, activities, objects, or situations)*?	0	1	2	3	4
8. Trouble remembering important parts of the stressful experience?	0	1	2	3	4
9. Having strong negative beliefs about yourself, other people, or the world *(for example, having thoughts such as: I am bad, there is something seriously wrong with me, no one can be trusted, the world is completely dangerous)*?	0	1	2	3	4
10. Blaming yourself or someone else for the stressful experience or what happened after it?	0	1	2	3	4
11. Having strong negative feelings such as fear, horror, anger, guilt, or shame?	0	1	2	3	4
12. Loss of interest in activities that you used to enjoy?	0	1	2	3	4
13. Feeling distant or cut off from other people?	0	1	2	3	4
14. Trouble experiencing positive feelings *(for example, being unable to feel happiness or have loving feelings for people close to you)*?	0	1	2	3	4
15. Irritable behavior, angry outbursts, or acting aggressively?	0	1	2	3	4
16. Taking too many risks or doing things that could cause you harm?	0	1	2	3	4
17. Being “superalert” or watchful or on guard?	0	1	2	3	4
18. Feeling jumpy or easily startled?	0	1	2	3	4
19. Having difficulty concentrating?	0	1	2	3	4
20. Trouble falling or staying asleep?	0	1	2	3	4

PCL Scoring

There are several ways in which to score the **PTSD Checklist (PCL).** Perhaps the easiest way to score the PCL is to add up all the items for a total severity score. A total score of 44 is considered to be PTSD positive for the general population while a total score of 50 is considered to be PTSD positive in military populations. A second way to score the PCL is to treat "moderately" or above (responses 3 through 5) as symptomatic and anything below "moderately" (1 and 2) as non-symptomatic. Then use the DSM scoring rules to make a diagnosis. That is:

- You need an endorsement of at least 1 B item (question #s 1-5)
- You need an endorsement of at least 3 C items (question #s 6-12)
- You need an endorsement of at least 2 D items (question #s 13-17)

However, please note that it is then possible to get a PTSD diagnosis with a total score of 29, which would be very low. It may therefore be best to use a combination of the two approaches. That is, the requisite number of items within each cluster are met at a 3 or above AND the total score is above the specified cut point.

Patient name: _____

Date of birth: _____

Alcohol screening questionnaire (AUDIT)

Drinking alcohol can affect your health and some medications you may take. Please help us provide you with the best medical care by answering the questions below.

One drink equals:  12 oz. beer  5 oz. wine  1.5 oz. liquor (one shot)

1. How often do you have a drink containing alcohol?	Never	Monthly or less	2 - 4 times a month	2 - 3 times a week	4 or more times a week
2. How many drinks containing alcohol do you have on a typical day when you are drinking?	0 - 2	3 or 4	5 or 6	7 - 9	10 or more
3. How often do you have five or more drinks on one occasion?	Never	Less than monthly	Monthly	Weekly	Daily or almost daily
4. How often during the last year have you found that you were not able to stop drinking once you had started?	Never	Less than monthly	Monthly	Weekly	Daily or almost daily
5. How often during the last year have you failed to do what was normally expected of you because of drinking?	Never	Less than monthly	Monthly	Weekly	Daily or almost daily
6. How often during the last year have you needed a first drink in the morning to get yourself going after a heavy drinking session?	Never	Less than monthly	Monthly	Weekly	Daily or almost daily
7. How often during the last year have you had a feeling of guilt or remorse after drinking?	Never	Less than monthly	Monthly	Weekly	Daily or almost daily
8. How often during the last year have you been unable to remember what happened the night before because of your drinking?	Never	Less than monthly	Monthly	Weekly	Daily or almost daily
9. Have you or someone else been injured because of your drinking?	No		Yes, but not in the last year		Yes, in the last year
10. Has a relative, friend, doctor, or other health care worker been concerned about your drinking or suggested you cut down?	No		Yes, but not in the last year		Yes, in the last year
	0	1	2	3	4

Have you ever been in treatment for an alcohol problem? ◯ Never ◯ Currently ◯ In the past

I	II	III	IV
0-3	4-9	10-13	14+

(For the Provider)

Scoring and interpreting the AUDIT:

1. Each response has a score ranging from 0 to 4. All response scores are added for a total score.

2. The total score correlates with a risk zone, which can be circled on the bottom left corner.

Score	Zone	Explanation	Action
0-3	I – Low Risk	"Someone using alcohol at this level is at low risk for health or social complications."	Positive Health Message – describe low risk drinking guidelines
4-9	II – Risky	"Someone using alcohol at this level may develop health problems or existing problems may worsen."	Brief intervention to reduce use
10-13	III – Harmful	"Someone using alcohol at this level has experienced negative effects from alcohol use."	Brief Intervention to reduce or abstain and specific follow-up appointment (Brief Treatment if available)
14+	IV – Severe	"Someone using alcohol at this level could benefit from more assessment and assistance."	Brief Intervention to accept referral to specialty treatment for a full assessment

Positive Health Message: An opportunity to educate patients about the NIAAA low-risk drinking levels and the risks of excessive alcohol use.

Brief Intervention to Reduce Use: Patient-centered discussion that uses Motivational Interviewing concepts to raise an individual's awareness of his/her substance use and enhance his/her motivation to change behavior. Brief interventions are typically 5-15 minutes, and should occur in the same session as the initial screening. Repeated sessions are more effective than a one-time intervention. The recommended behavior change is to cut back to low-risk drinking levels unless there are other medical reasons to abstain (liver damage, pregnancy, medication contraindications, etc.).

Brief Intervention to Reduce or Abstain (Brief Treatment if available) & Follow-up: Patients with numerous or serious negative consequences from their alcohol use, or patients who likely have an alcohol use disorder who cannot or are not interested in obtaining specialized treatment, should receive more numerous and intensive BIs with follow up. The recommended behavior change is to cut back to low-risk drinking levels or abstain from use. Brief treatment is 1 to 5 sessions, each 15-60 minutes. Refer for brief treatment if available. If brief treatment is not available, secure follow-up in 2-4 weeks.

Brief Intervention to Accept Referral: The focus of the brief intervention is to enhance motivation for the patient to accept a referral to specialty treatment. If accepted, the provider should use a proactive process to facilitate access to specialty substance use disorder treatment for diagnostic assessment and, if warranted, treatment. The recommended behavior change is to abstain from use and accept the referral.

More resources: www.sbirtoregon.org

***** Johnson J, Lee A, Vinson D, Seale P. "Use of AUDIT-Based Measures to Identify Unhealthy Alcohol Use and Alcohol Dependence in Primary Care: A Validation Study." Alcohol Clin Exp Res, Vol 37, No S1, 2013: pp E253–E259

Drug Abuse Screening Test, DAST-10

The following questions concern information about your possible involvement with drugs *not including alcoholic beverages* during the past 12 months.

"Drug abuse" refers to (1) the use of prescribed or over-the-counter drugs in excess of the directions, and (2) any nonmedical use of drugs.

The various classes of drugs may include cannabis (marijuana, hashish), solvents (e.g., paint thinner), tranquilizers (e.g., Valium), barbiturates, cocaine, stimulants (e.g., speed), hallucinogens (e.g., LSD) or narcotics (e.g., heroin). Remember that the questions *do not* include alcoholic beverages.

Please answer every question. If you have difficulty with a statement, then choose the response that is mostly right.

In the past 12 months…		**Circle**	
1.	Have you used drugs other than those required for medical reasons?	Yes	No
2.	Do you abuse more than one drug at a time?	Yes	No
3.	Are you unable to stop abusing drugs when you want to?	Yes	No
4.	Have you ever had blackouts or flashbacks as a result of drug use?	Yes	No
5.	Do you ever feel bad or guilty about your drug use?	Yes	No
6.	Does your spouse (or parents) ever complain about your involvement with drugs?	Yes	No
7.	Have you neglected your family because of your use of drugs?	Yes	No
8.	Have you engaged in illegal activities in order to obtain drugs?	Yes	No
9.	Have you ever experienced withdrawal symptoms (felt sick) when you stopped taking drugs?	Yes	No
10.	Have you had medical problems as a result of your drug use (e.g. memory loss, hepatitis, convulsions, bleeding)?	Yes	No
Scoring: Score 1 point for each question answered "Yes," except for question 3 for which a "No" receives 1 point.		**Score:**	

Interpretation of Score		
Score	**Degree of Problems Related to Drug Abuse**	**Suggested Action**
0	No problems reported	None at this time
1-2	Low level	Monitor, re-assess at a later date
3-5	Moderate level	Further investigation
6-8	Substantial level	Intensive assessment
9-10	Severe level	Intensive assessment

Drug Abuse Screening Test (DAST-10). (Copyright 1982 by the Addiction Research Foundation.)

GAD-7 Anxiety

Over the last two weeks, how often have you been bothered by the following problems?	Not at all	Several days	More than half the days	Nearly every day
1. Feeling nervous, anxious, or on edge	0	1	2	3
2. Not being able to stop or control worrying	0	1	2	3
3. Worrying too much about different things	0	1	2	3
4. Trouble relaxing	0	1	2	3
5. Being so restless that it is hard to sit still	0	1	2	3
6. Becoming easily annoyed or irritable	0	1	2	3
7. Feeling afraid, as if something awful might happen	0	1	2	3

Column totals _____ + _____ + _____ + _____ =

Total score _____

If you checked any problems, how difficult have they made it for you to do your work, take care of things at home, or get along with other people?

Not difficult at all	Somewhat difficult	Very difficult	Extremely difficult
☐	☐	☐	☐

Source: Primary Care Evaluation of Mental Disorders Patient Health Questionnaire (PRIME-MD-PHQ). The PHQ was developed by Drs. Robert L. Spitzer, Janet B.W. Williams, Kurt Kroenke, and colleagues. For research information, contact Dr. Spitzer at ris8@columbia.edu. PRIME-MD® is a trademark of Pfizer Inc. Copyright© 1999 Pfizer Inc. All rights reserved. Reproduced with permission

Scoring GAD-7 Anxiety Severity

This is calculated by assigning scores of 0, 1, 2, and 3 to the response categories, respectively, of “not at all,” “several days,” “more than half the days,” and “nearly every day.”
GAD-7 total score for the seven items ranges from 0 to 21.

0–4: minimal anxiety

5–9: mild anxiety

10–14: moderate anxiety

15–21: severe anxiety

PATIENT HEALTH QUESTIONNAIRE-9 (PHQ-9)

Over the last 2 weeks, how often have you been bothered by any of the following problems? *(Use "✔" to indicate your answer)*	**Not at all**	**Several days**	**More than half the days**	**Nearly every day**
1. Little interest or pleasure in doing things	0	1	2	3
2. Feeling down, depressed, or hopeless	0	1	2	3
3. Trouble falling or staying asleep, or sleeping too much	0	1	2	3
4. Feeling tired or having little energy	0	1	2	3
5. Poor appetite or overeating	0	1	2	3
6. Feeling bad about yourself — or that you are a failure or have let yourself or your family down	0	1	2	3
7. Trouble concentrating on things, such as reading the newspaper or watching television	0	1	2	3
8. Moving or speaking so slowly that other people could have noticed? Or the opposite — being so fidgety or restless that you have been moving around a lot more than usual	0	1	2	3
9. Thoughts that you would be better off dead or of hurting yourself in some way	0	1	2	3

FOR OFFICE CODING __0__ **+** _____ **+** _____ **+** _____

=Total Score: _____

If you checked off any problems, how difficult have these problems made it for you to do your work, take care of things at home, or get along with other people?

Not difficult at all	**Somewhat difficult**	**Very difficult**	**Extremely difficult**
☐	☐	☐	☐

Developed by Drs. Robert L. Spitzer, Janet B.W. Williams, Kurt Kroenke and colleagues, with an educational grant from Pfizer Inc. No permission required to reproduce, translate, display or distribute.